

RATIONALITY OF W -ALGEBRAS, I: BERSHADSKY-POLYAKOV VERTEX ALGEBRAS

TOMOYUKI ARAKAWA

ABSTRACT. We prove the conjecture of Kac-Wakimoto on the rationality of exceptional W -algebras for the first non-trivial series, namely, for the Bershadsky-Polyakov vertex algebras $W_3^{(2)}$ at level $k = p/2 - 3$ with $p = 3, 5, 7, \dots$. This gives new examples of rational conformal field theories.

1. INTRODUCTION

Recently, a remarkable family of W -algebras associated with simple Lie algebras and their *non-principal* nilpotent elements, called *exceptional W -algebras*, has been discovered by Kac and Wakimoto [KW08]. In [KW08] it was conjectured that with an exceptional W -algebra one can associated a rational conformal field theory.

As a first step to resolve the Kac-Wakimoto conjecture we have proved in the previous article [Ara10b] that (non-principal) exceptional W -algebras are *lisse*, or equivalently [Ara10a], C_2 -cofinite. Therefore it remains [Zhu96, Hua08] to show that exceptional W -algebras are *rational*, i.e., that the representations are completely reducible, in order to prove the conjecture. In this article we prove the rationality of the first non-trivial series of exceptional W -algebras, that is, the *Bershadsky-Polyakov (vertex) algebras* $W_3^{(2)}$ [Pol90, Ber91] at level $k = p/2 - 3$ with $p = 3, 5, 7, \dots$. The vertex algebra $W_3^{(2)}$ is the W -algebra associated with $\mathfrak{g} = \mathfrak{sl}_3$ and its minimal nilpotent element.

Let us state our main result more precisely: Let $\mathcal{W}_k$ denote the unique simple quotient of $W_3^{(2)}$ at level $k \neq -3$.

Main Theorem (Conjectured by Kac and Wakimoto [KW08]). *Let $k = p/2 - 3$ with $p = 3, 5, 7, \dots$. Then the vertex algebra $\mathcal{W}_k$ is rational. The simple $\mathcal{W}_k$ -modules are parameterized by the set of integral dominant weights of $\widehat{\mathfrak{sl}}_3$ of level $p - 3$. These simple modules can be obtained by the quantum BRST reduction from irreducible admissible representations of $\widehat{\mathfrak{sl}}_3$ of level k .*

For $p = 3$, $\mathcal{W}_{3/2-3}$ is one-dimensional. In the remaining cases $\mathcal{W}_{p/2-3}$ are conformal with negative central charges. They have features in common with the $\widehat{\mathfrak{sl}}_2$ -integrable affine vertex algebras in the sense that their Zhu algebras are closely related with Smith's algebra [Smi90] and that the following relations hold:

$$: G^+(z)^{p-2} :=: G^-(z)^{p-2} := 0.$$

This work is partially supported by the JSPS Grant-in-Aid for Scientific Research (B) No. 20340007.

2. BERSHADSKY-POLYAKOV ALGEBRAS AT EXCEPTIONAL LEVELS.

Let $\mathcal{W}^k$ denote the Bershadsky-Polyakov (vertex) algebra $W_3^{(2)}$ at level $k \neq -3$, which is the vertex algebra freely generated by the fields $J(z), G^\pm(z), T(z)$ with the following OPE's:

$$\begin{aligned}
J(z)J(w) &\sim \frac{2k+3}{3(z-w)^2}, \quad G^\pm(z)G^\pm(w) \sim 0, \\
J(z)G^\pm(w) &\sim \pm \frac{1}{z-w}G^\pm(w), \\
T(z)T(w) &\sim -\frac{(2k+3)(3k+1)}{2(k+3)(z-w)^4} + \frac{2}{(z-w)^2}T(w) + \frac{1}{z-w}\partial T(w), \\
T(z)G^\pm(w) &\sim \frac{3}{2(z-w)^2}G^\pm(w) + \frac{1}{z-w}\partial G^\pm(w), \\
T(z)J(w) &\sim \frac{1}{(z-w)^2}J(w) + \frac{1}{z-w}\partial J(w), \\
G^+(z)G^-(w) &\sim \frac{(k+1)(2k+3)}{(z-w)^3} + \frac{3(k+1)}{(z-w)^2}J(w) \\
&\quad + \frac{1}{z-w} \left(3 : J(w)^2 : + \frac{3(k+1)}{2}\partial J(w) - (k+3)T(w) \right).
\end{aligned}$$

As in introduction we denote by $\mathcal{W}_k$ the unique simple quotient of $\mathcal{W}^k$.

Theorem 2.0.1 ([Ara10b]). *Let k, p be as in Main Theorem. Then $\mathcal{W}_k$ is lisse.*

Set

$$L(z) = \sum_{n \in \mathbb{Z}} L_n z^{-n-2} = T(z) + \frac{1}{2}\partial J(w).$$

This defines a conformal vector of $\mathcal{W}^k$ with central charge

$$c(k) = -\frac{4(k+1)(2k+3)}{k+3} = -\frac{4(p-4)(p-3)}{p},$$

which gives J, G^+, G^- conformal weights 1, 1, and 2, respectively. Hence $\mathcal{W}^k$ is $\mathbb{Z}_{\geq 0}$ -graded with respect to the Hamiltonian L_0 . We expand the corresponding fields accordingly:

$$J(z) = \sum_{n \in \mathbb{Z}} J_n z^{-n-1}, \quad G^+(z) = \sum_{n \in \mathbb{Z}} G_n^+ z^{-n-1}, \quad G^-(z) = \sum_{n \in \mathbb{Z}} G_n^- z^{-n-2}.$$

Then we have

$$\begin{aligned}
[J_m, J_n] &= \frac{2k+3}{3}m\delta_{m+n,0}, \quad [J_m, G_n] = G_{m+n}, \quad [J_m, F_n] = -F_{m+n}, \\
[L_m, J_n] &= -nJ_{m+n} - \frac{(2k+3)(m+1)m}{6}\delta_{m+n,0}, \\
[L_m, G_n^+] &= -nG_{m+n}^+, \quad [L_m, G_n^-] = (m-n)G_{m+n}^-, \\
[G_m^+, G_n^-] &= 3(J^2)_{m+n} + (3(k+1)m - (2k+3)(m+n+1))J_{m+n} - (k+3)L_{m+n} \\
&\quad + \frac{(k+1)(2k+3)m(m+1)}{2}\delta_{m+n,0},
\end{aligned}$$

where $\sum_{n \in \mathbb{Z}} (J^2)_n z^{-n-2} := : J(z)^2 :$.

For $(\xi, \chi) \in \mathbb{C}^2$, let $L(\xi, \chi)$ be the irreducible representation of $\mathcal{W}^k$ generated by the vector $|\xi, \chi\rangle$ such that

$$\begin{aligned} J_0|\xi, \chi\rangle &= \xi|\xi, \chi\rangle, & J_n|\xi, \chi\rangle &= 0 \quad (n > 0), \\ L_0|\xi, \chi\rangle &= \chi|\xi, \chi\rangle, & L_n|\xi, \chi\rangle &= 0 \quad (n > 0), \\ G_n^-|\xi, \chi\rangle &= 0 \quad (n \geq 0), & G_n^+|\xi, \chi\rangle &= 0 \quad (n \geq 1). \end{aligned}$$

By Theorem 2.0.1, any simple $\mathcal{W}_k$ -module is of the form $L(\xi, \lambda)$ with some ξ and χ . (It is important that the lisse condition is defined independent of the grading.)

For a $\mathcal{W}^k$ -module M set $M_{a,d} = \{m \in M; J_0m = am, L_0m = dm\}$. It is clear that $L(\xi, \chi) = \bigoplus_{\substack{(a,d) \in \mathbb{C}^2 \\ d \in \chi + \mathbb{Z}_{\geq 0}}} L(\xi, \chi)_{a,d}$, $\dim L(\xi, \chi)_{\xi, \chi} = 1$. Let $L(\xi, \chi)_{\text{top}} = \{v \in$

$L(\xi, \chi); L_0v = \chi v\} = \bigoplus_a L(\xi, \chi)_{a, \chi}$. By definition $L(\xi, \chi)_{\text{top}}$ is spanned by the vectors $(G_0^+)^i|\xi, \chi\rangle$ with $i \geq 0$.

Following [Smi90] set

$$g(\xi, \chi) = -(3\xi^2 - (2k+3)\xi - (k+3)\chi),$$

so that $G_0^-G_0^+|\xi, \chi\rangle = g(\xi, \chi)|\xi, \chi\rangle$. We have

$$G_0^-(G_0^+)^i|\xi, \chi\rangle = ih_i(\xi, \chi)(G_0^+)^{i-1}|\xi, \chi\rangle,$$

where

$$\begin{aligned} h_i(\xi, \chi) &= \frac{1}{i}(g(\xi, \chi) + g(\xi+1, \chi) + \cdots + g(\xi+i-1, \chi)) \\ &= -i^2 + ki - 3\xi i + 3i - 3\xi^2 - k + 2k\xi + 6\xi + k\chi + 3\chi - 2. \end{aligned}$$

Hence we have the following assertion.

Proposition 2.0.2. *The space $L(\xi, \chi)_{\text{top}}$ is n -dimensional only if $h_n(\xi, \chi) = 0$.*

Define

$$\Delta(-J, z) = z^{-J_0} \exp\left(\sum_{k=1}^{\infty} (-1)^{k+1} \frac{-J_k}{kz^k}\right),$$

and set

$$\sum_{n \in \mathbb{Z}} \psi(a_{(n)}) z^{-n-1} = Y(\Delta(-J, z)a, z)$$

for $a \in \mathcal{W}^k$. Then for any $\mathcal{W}^k$ -module M , we can define on M a new $\mathcal{W}^K$ -module structure by twisting the action of $\mathcal{W}^k$ as $a_{(n)} \mapsto \psi(a_{(n)})$ ([Li97]). We denote by $\psi(M)$ thus obtained $\mathcal{W}^k$ -module from M .

Proposition 2.0.3. *Suppose that $\dim L(\xi, \chi)_{\text{top}} = i$. Then*

$$\psi(L(\xi, \chi)) \cong L\left(\xi + i - 1 - \frac{2k+3}{3}, \chi - (\xi - i + 1) + \frac{2k+3}{3}\right).$$

Proof. The assertion follows from the fact that

$$\begin{aligned} \psi(J_n) &= J_n - \frac{2k+3}{3}\delta_{n,0}, & \psi(L_n) &= L_n - J_n + \frac{2k+3}{3}, \\ \psi(G_n^+) &= G_{n-1}^+, & \psi(G_n^-) &= G_{n+1}^-. \end{aligned}$$

□

By solving the equation

$$h_i(\xi, \chi) = h_j(\xi + i - 1 - \frac{2k+3}{3}, \chi - (\xi - i + 1) + \frac{2k+3}{3})$$

we obtain the following assertion.

Proposition 2.0.4. *Suppose that $\dim L(\xi, \chi)_{\text{top}} = i$ and $\dim \psi(L(\xi, \chi))_{\text{top}} = j$. Then*

$$\begin{aligned} \xi &= \xi_{i,j} := \frac{1}{3}(-2i - j + 2k + 6), \\ \chi &= \chi_{i,j} := \frac{i^2 + ji - ki - 3i + j^2 - 6j - 2jk + 3k + 6}{3(k+3)}. \end{aligned}$$

Proposition 2.0.5. *Let k, p be as in Main Theorem. Then $(G_{-1}^+)^{p-2}\mathbf{1}$ belongs to the maximal ideal of $\mathcal{W}^k$.*

Proof. Because $\xi_{1,p-2} = \chi_{1,p-2} = 0$, the corresponding $\mathbf{1} \mapsto |\xi_{1,p-2}, \chi_{1,p-2}\rangle$ gives an isomorphism $\mathcal{W}_k \cong L(\xi_{1,p-2}, \chi_{1,p-2})$. Because $h_{p-2}(\xi_{1,p-2} - (2k+3)/2, \chi_{1,p-2} + (2k+3)/3) = 0$, from Proposition 2.0.3 it follows that $\psi(\mathcal{W}_k)_{\text{top}}$ is at most $p-2$ -dimensional. Hence $(G_{-1}^+)^{p-2}\mathbf{1} = 0$. $\square$

Remark 2.0.6. One can show that in fact $(G_{-1}^+)^{p-2}$ generates the maximal ideal of $\mathcal{W}^k$. However we do not need this fact.

Proposition 2.0.7. *Let k, p be as in Main Theorem. Then any simple $\mathcal{W}_k$ -module is isomorphic to $L(\xi_{i,j}, \chi_{i,j})$ for some (i, j) such that $1 \leq i \leq p-2$, $1 \leq j \leq p-i-1$.*

Proof. Let $L(\xi, \chi)$ be a simple $\mathcal{W}_k$ -module. Because $G^+(z)^{p-2} := 0$ by Proposition 2.0.5, $L(\xi, \chi)_{\text{top}}$ is at most $(p-2)$ -dimensional. Since $\psi(L(\xi, \chi))$ is also a $\mathcal{W}_k$ -module we have $(\xi, \chi) = (\xi_{i,j}, \chi_{i,j})$ for some $1 \leq i, j \leq p-2$. Because $\psi(\psi(L(\xi_{i,j}, \chi_{i,j})))$ is also a $\mathcal{W}_k$ -module it follows that $\xi_{i,j} + i - 1 - \frac{2k+3}{3} = \frac{i-j}{3} \leq \frac{-2j-1+2k+6}{3} = \frac{p-2j-1}{3}$. Hence $j \leq p-i-1$. $\square$

Remark 2.0.8. The simple $\mathcal{W}^k$ -modules $L(\xi_{i,j}, \chi_{i,j})$ with $1 \leq i \leq p-2$, $1 \leq j \leq p-i-1$, are mutually non-isomorphic because their highest weights are distinct.

3. PROOF OF MAIN THEOREM

Let k, p be as in Main Theorem.

Let $\mathfrak{g} = \mathfrak{sl}_3$ as in introduction, $\mathfrak{h} \subset \mathfrak{g}$ be the Cartan subalgebra of $\mathfrak{g}$ consisting of diagonal matrixes. Set $h_i = E_{i,i} - E_{i+1,i+1}$, $h_\theta = h_1 + h_2$, $e_i = e_{\alpha_i} = E_{i,i+1}$, $f_i = f_{\alpha_i} = E_{i+1,i}$ for $i = 1, 2$, $e_\theta = E_{1,3}$, $f_\theta = E_{3,1}$, where $E_{i,j}$ is the matrix element. We equip $\mathfrak{g}$ the invariant form $(x|y) = \text{tr}(xy)$. Set $\bar{\Lambda}_1 = (2h_1 + h_2)/3$, $\bar{\Lambda}_1 = (h_1 + 2h_2)/3$, so that $(\bar{\Lambda}_i|h_j) = \delta_{i,j}$.

Let $\hat{\mathfrak{g}} = \mathfrak{g}[t, t^{-1}] \oplus \mathbb{C}K \oplus \mathbb{C}D$ be the (non-twisted) affine Kac-Moody algebra associated with $\mathfrak{g}$, where K is the central element and D is the degree operator. Let $\hat{\mathfrak{h}} = \mathfrak{h} \oplus \mathbb{C}K \oplus \mathbb{C}D \subset \hat{\mathfrak{g}}$ the standard Cartan subalgebra, $\hat{\mathfrak{h}}^* = \mathfrak{h}^* \oplus \mathbb{C}\Lambda_0 \oplus \mathbb{C}\delta$ the dual of $\hat{\mathfrak{h}}$, where Λ_0 and δ are elements dual to K and D , respectively.

Let us consider the minimal nilpotent element f_θ and the corresponding Dynkin grading of $\mathfrak{g}$: $\mathfrak{g}_j = \{u \in \mathfrak{g}; [h_\theta, u] = 2ju\}$. Denote by $H_{f_\theta}^{\infty+0}(\cdot)$ the BRST cohomology of the generalized quantized Drinfeld-Sokolov reduction associated with $(\mathfrak{g}, f_\theta)$

and the corresponding Dynkin grading. Then we have [KRW03, KW04] the vertex algebra isomorphism

$$\mathcal{W}^k \simeq H_{f_\theta}^{\frac{\infty}{2}+0}(V^k(\mathfrak{g})),$$

which is given by the following assignment:

$$\begin{aligned} J(z) &\mapsto J^{-\bar{\Lambda}_1+\bar{\Lambda}_2}(z) - : \Phi_1(z)\Phi_2(z) :, \\ G^+(z) &\mapsto J^{f_1}(z) - : J^{h_1}(z)\Phi_2(z) : + : \Phi_1(z)\Phi_2(z)^2 : - (k+1)\partial\Phi_2(z), \\ G^-(z) &\mapsto -J^{f_2}(z) - : J^{h_2}(z)\Phi_1(z) : - : \Phi_1(z)^2\Phi_2(z) : - (k+1)\partial\Phi_1(z), \end{aligned}$$

Here $J^u(z) = u(z) - \sum_{\beta, \gamma \in \{\alpha_1, \alpha_2, \theta\}} c_{u, f_\beta}^{f_\gamma} : \psi_\beta^*(z)\psi_\gamma(z) :$ for $u \in \mathfrak{g}$, $c_{u_1, u_2}^{u_3}$ is the structure constant, $\psi_\alpha(z)$, $\psi_\alpha^*(z)$ with $\alpha \in \{\alpha_1, \alpha_2, \theta\}$ are fermionic ghosts satisfying

$$(1) \quad \psi_\alpha(z)\psi_\beta^*(w) \sim \frac{\delta_{\alpha, \beta}}{z-w}, \quad \psi_\alpha(z)\psi_\beta(w) \sim \psi_\alpha^*(z)\psi_\beta^*(w) \sim 0,$$

$\Phi_1(z)$, $\Phi_2(z)$ are bosonic ghosts satisfying

$$\Phi_1(z)\Phi_2(w) \sim \frac{1}{z-w}, \quad \Phi_i(z)\Phi_i(w) \sim 0,$$

and the BRST differential is the zero mode of the field

$$\begin{aligned} Q(z) &= \sum_{\alpha \in \{\alpha_1, \alpha_2, \theta\}} e_\alpha(z)\psi_\alpha^*(w) - : \psi_{\alpha_1}^*(z)\psi_{\alpha_2}^*(z)\psi_\theta(z) : \\ &\quad + \Phi_1(z)\psi_{\alpha_1}^*(z) + \Phi_2(z)\psi_{\alpha_2}(z) + \psi_\theta(z), \end{aligned}$$

Let $\mathcal{O}_k$ be the category $\mathcal{O}$ of $\widehat{\mathfrak{g}}$ at level k , $\mathbf{L}_\lambda$ the irreducible representation of $\widehat{\mathfrak{g}}$ with highest weight λ . Let $\mathcal{W}^k\text{-Mod}$ be the category of $\mathcal{W}^k$ -modules.

Theorem 3.0.9 ([Ara05]). (i) *The functor $H_{f_\theta}^{\frac{\infty}{2}+0}(?) : \mathcal{O}_k \rightarrow \mathcal{W}^k\text{-Mod}$ is exact.*

(ii) *For $\mathbf{L}_\lambda \in \mathcal{O}_k$ we have $H_{f_\theta}^{\frac{\infty}{2}+0}(\mathbf{L}_\lambda) = 0$ if and only if $\lambda(\alpha_0^\vee) \in \{0, 1, 2, \dots\}$. Otherwise $H_{f_\theta}^{\frac{\infty}{2}+0}(\mathbf{L}_\lambda)$ is irreducible.*

Let Adm^k be the set of admissible weights [KW89] of $\widehat{\mathfrak{g}}$ of level k , and put

$$\text{Adm}_+^k = \{\lambda \in \text{Adm}^k; \bar{\lambda} \text{ is an integral dominant weight of } \mathfrak{g}\},$$

where $\widehat{\mathfrak{h}}^* \ni \lambda \mapsto \bar{\lambda} \in \mathfrak{h}^*$ is the restriction. Then

$$\text{Adm}_+^k = \{\bar{\mu} + k\Lambda_0; \mu \in \widehat{P}_{++}^{p-3}\},$$

where $\widehat{P}_{++}^{p-3}$ is the set of integral dominant weights of $\widehat{\mathfrak{g}}$ of level $p-3$. Explicitly, we have

$$\text{Adm}_+^k = \{\lambda_{i,j}; 1 \leq i \leq p-2, 1 \leq j \leq p-i-1\},$$

where

$$\lambda_{i,j} = (i-1)\bar{\Lambda}_1 + (p-i-j-1)\bar{\Lambda}_2 + k\Lambda_0.$$

Note that

$$(2) \quad \xi_{i,j} = (\lambda_{i,j} | -\bar{\Lambda}_1 + \bar{\Lambda}_2), \quad \chi_{i,j} = \frac{(\lambda_{i,j} | \lambda_{i,j} + 2\bar{\rho})}{2(k+3)} - (\lambda_{i,j} | \bar{\Lambda}_2),$$

where $\bar{\rho} = \bar{\Lambda}_1 + \bar{\Lambda}_2$.

Recall the following result of Malikov and Frenkel [MF99].

Theorem 3.0.10 ([MF99]). *For $\lambda \in \text{Adm}_+^k$, $\mathbf{L}_\lambda$ is a module over $\mathbf{L}_{k\Lambda_0}$.*

Proposition 3.0.11. *For $\lambda_{i,j} \in \text{Adm}_+^k$, $H_{f_\theta}^{\frac{\infty}{2}+0}(\mathbf{L}_{\lambda_{i,j}})$ is a simple $\mathcal{W}_k$ -module isomorphic to $L(\xi_{i,j}, \chi_{i,j})$.*

Proof. By Theorem 3.0.9 we have $\mathcal{W}_k \cong H_{f_\theta}^{\frac{\infty}{2}+0}(\mathbf{L}_{k\Lambda_0})$. Hence by the functoriality of $H_{f_\theta}^{\frac{\infty}{2}+0}(?)$, Theorem 3.0.10 immediately gives that $H_{f_\theta}^{\frac{\infty}{2}+0}(\mathbf{L}_{\lambda_{i,j}})$ is a $\mathcal{W}_k$ -module. This module is (nonzero and) irreducible by Theorem 3.0.9. Let v be the image of the highest weight vector of $\mathbf{L}_{\lambda_{i,j}}$ in $H_{f_\theta}^{\frac{\infty}{2}+0}(\mathbf{L}_{\lambda_{i,j}})$. By (2) and the fact [KW04] that the image of $L(z)$ in $\mathcal{W}^k$ is cohomologous to

$$L_{\mathfrak{g}}(z) + L_{\text{ch}}(z) + L_{\Phi}(z) + \partial J^{\bar{\Lambda}_2}(z),$$

where $L_{\mathfrak{g}}(z)$ is the Sugawara operator of $\mathfrak{g}$, $L_{\text{ch}}(z) = -\sum_{\alpha=\alpha_1, \alpha_2, \theta} \phi_\alpha(z) \partial \phi_\alpha^*(z)$, $L_{\Phi}(z) = \frac{1}{2}(\Phi_2(z) \partial \Phi_1(z) - \partial \Phi_1(z) \Phi_2(z))$, it is straightforward to check that the assignment $|\xi_{i,j}, \chi_{i,j}\rangle \mapsto v$ gives a $\mathcal{W}^k$ -module homomorphism. By the irreducibility, this must be an isomorphism. $\square$

By Propositions 2.0.7 and 3.0.11, $\{H_{f_\theta}^{\frac{\infty}{2}+0}(\mathbf{L}_\lambda); \lambda \in \text{Adm}_+^k\}$ gives the complete set of isomorphism classes of simple $\mathcal{W}_k$ -modules. Therefore Main Theorem now follows immediately from the following important result of Gorelik and Kac [GK09].

Theorem 3.0.12 ([GK09, Corollary 8.8.9]). *For any $\lambda, \mu \in \text{Adm}^k$, we have*

$$\text{Ext}_{\mathcal{W}^k\text{-Mod}}^1(H_{f_\theta}^{\frac{\infty}{2}+0}(\mathbf{L}_\lambda), H_{f_\theta}^{\frac{\infty}{2}+0}(\mathbf{L}_\mu)) = 0.$$

REFERENCES

- [Ara05] Tomoyuki Arakawa. Representation theory of superconformal algebras and the Kac-Roan-Wakimoto conjecture. *Duke Math. J.*, Vol. 130, No. 3, pp. 435–478, 2005.
- [Ara10a] T. Arakawa. A remark on the C_2 -cofiniteness condition on vertex algebras. *ArXiv e-prints*, April 2010.
- [Ara10b] T. Arakawa. Associated varieties of modules over Kac-Moody algebras and C_2 -cofiniteness of W -algebras. *ArXiv e-prints*, April 2010.
- [Ber91] Michael Bershadsky. Conformal field theories via Hamiltonian reduction. *Comm. Math. Phys.*, Vol. 139, No. 1, pp. 71–82, 1991.
- [GK09] M. Gorelik and V. Kac. On complete reducibility for infinite-dimensional Lie algebras. *ArXiv e-prints*, May 2009.
- [Hua08] Yi-Zhi Huang. Vertex operator algebras and the Verlinde conjecture. *Commun. Contemp. Math.*, Vol. 10, No. 1, pp. 103–154, 2008.
- [KRW03] Victor Kac, Shi-Shyr Roan, and Minoru Wakimoto. Quantum reduction for affine superalgebras. *Comm. Math. Phys.*, Vol. 241, No. 2-3, pp. 307–342, 2003.
- [KW89] V. G. Kac and M. Wakimoto. Classification of modular invariant representations of affine algebras. In *Infinite-dimensional Lie algebras and groups (Luminy-Marseille, 1988)*, Vol. 7 of *Adv. Ser. Math. Phys.*, pp. 138–177. World Sci. Publ., Teaneck, NJ, 1989.
- [KW04] Victor G. Kac and Minoru Wakimoto. Quantum reduction and representation theory of superconformal algebras. *Adv. Math.*, Vol. 185, No. 2, pp. 400–458, 2004.
- [KW08] Victor G. Kac and Minoru Wakimoto. On rationality of W -algebras. *Transform. Groups*, Vol. 13, No. 3-4, pp. 671–713, 2008.
- [Li97] Haisheng Li. The physics superselection principle in vertex operator algebra theory. *J. Algebra*, Vol. 196, No. 2, pp. 436–457, 1997.
- [MF99] F. G. Malikov and I. B. Frenkel'. Annihilating ideals and tilting functors. *Funktsional. Anal. i Prilozhen.*, Vol. 33, No. 2, pp. 31–42, 95, 1999.

- [Pol90] A. M. Polyakov. Gauge transformations and diffeomorphisms. *Internat. J. Modern Phys. A*, Vol. 5, No. 5, pp. 833–842, 1990.
- [Smi90] S. P. Smith. A class of algebras similar to the enveloping algebra of $\mathfrak{sl}(2)$. *Trans. Amer. Math. Soc.*, Vol. 322, No. 1, pp. 285–314, 1990.
- [Zhu96] Yongchang Zhu. Modular invariance of characters of vertex operator algebras. *J. Amer. Math. Soc.*, Vol. 9, No. 1, pp. 237–302, 1996.

DEPARTMENT OF MATHEMATICS, NARA WOMEN'S UNIVERSITY, NARA 630-8506, JAPAN
E-mail address: `arakawa@cc.nara-wu.ac.jp`